## **Techniques of integration:** Substitution

**Observation:** If u = g(x), then du = g'(x) dx.

We can use this observation to simplify integrals that have the form

$$\int f(g(x)) \, g'(x) \, dx,$$

by **substituting** u = g(x), which entails du = g'(x) dx:

$$\int f(\widetilde{g(x)}) \, \widetilde{g'(x) \, dx} = \int f(u) \, du$$

**Example 1:** The substitution  $u = x^2 + 3$ , du = 2x dx gives

$$\int 2x\sqrt{x^2+3} \, dx = \int (\overbrace{x^2+3}^u)^{1/2} \overbrace{2x \, dx}^{du}$$
$$= \int u^{1/2} \, du = \frac{u^{3/2}}{3/2} + C = \frac{2}{3}u^{3/2} + C$$
$$= \frac{2}{3}(x^2+3)^{3/2} + C$$

(\*) The key is *recognizing* that a substitution is possible.

(\*) What we are looking for in the integrand is a *product*, with one factor being a *composite function*, e.g., f(g(x)), and the other factor being the *derivative of the argument* of the first factor, i.e., g'(x).

Example 2: Compute

$$\int \frac{2x}{x^2 + 4} \, dx$$

The integrand looks like a quotient, but really it's a product, and one of the factors is composite:

$$\int \frac{2x}{x^2 + 4} \, dx = \int (x^2 + 4)^{-1} \, 2x \, dx$$

Substituting  $u = x^2 + 4$  and du = 2x dx makes the integral easy to compute:

$$\int (\overbrace{x^2+4}^{u})^{-1} \underbrace{2x \, dx}^{du} = \int u^{-1} \, du = \ln|u| + C = \ln(x^2+4) + C.$$

**Observation:** We can multiply or divide any equation by a *nonzero* constant. This makes substitution more flexible.

Example 3: Compute

$$\int 5x^2 \sqrt[4]{2x^3 + 1} \, dx$$

The natural choice for a substitution is  $u = 2x^3 + 1$ , which means that  $du = 6x^2 dx$ . Dividing this relation by 6 shows that

$$x^{2} dx = \frac{1}{6} du \implies 5x^{2} dx = \frac{5}{6} du,$$

so the substitution  $u = 2x^3 + 1$  will work here:

$$\int 5x^2 \sqrt[4]{2x^3 + 1} \, dx = \int (2x^3 + 1)^{1/4} \underbrace{5x^2 \, dx}_{5x^2 \, dx}$$
$$= \frac{5}{6} \int u^{1/4} \, du = \frac{2}{3} u^{5/4} + C$$
$$= \frac{2}{3} (2x^3 + 1)^{5/4} + C$$

**Observation:** If u = ax + b, then du = a dx so that  $dx = \frac{1}{a} du$ . This means that we can always make the following simplification

$$\int f(ax+b) \, dx = \frac{1}{a} \int f(u) \, du$$

by substituting u = ax + b.

Example 4: Compute

$$\int \sqrt[3]{5x+2} \, dx = \int (5x+2)^{1/3} \, dx.$$

The substitution u = 5x + 2,  $du = 5 dx \implies dx = \frac{1}{5} du$  will work here:

$$\int \underbrace{(5x+2)^{1/3} \cdot \frac{1}{5}^{du}}_{dx} = \int u^{1/3} \cdot \frac{1}{5} \, du = \frac{3}{20} u^{4/3} + C = \frac{3}{20} (5x+2)^{4/3} + C$$

**Example 5:** Find the demand equation for a firm whose marginal revenue function is

$$\frac{dr}{dq} = 10\sqrt[3]{1000 - 0.5q}.$$

**Observation:** Since r = pq, by definition, we can find the demand equation for this firm by

(i) Finding the revenue function r(q), and

(ii) Writing p = r(q)/q.

**Step 1.** Integrate, by substituting u = 1000 - 0.5q,  $du = -0.5 dq \implies dq = -2 du$ :

$$\int 10\sqrt[3]{1000 - 0.5q} \, dq = 10 \int (\underbrace{1000 - 0.5q}^{u})^{1/3} \underbrace{\frac{-2 \, du}{dq}}_{= -20 \int u^{1/3} \, du}_{= -15u^{4/3} + C} = -15(1000 - 0.5q)^{4/3} + C$$

Conclusion 1:  $r = C - 15(1000 - 0.5q)^{4/3}$ 

Step 2: Solve for C:

$$0 = r(0) = C - 15(1000)^{4/3} \implies C = 150000.$$

Conclusion:  $r = 150000 - 15(1000 - 0.5q)^{4/3}$ .

**Step 3:** Write down the demand equation:

$$p = \frac{r}{q} = \frac{15\left(10000 - (1000 - 0.5q)^{4/3}\right)}{q}$$

**Observation:** Linear equations are easy to solve. For example, if u = ax + b, then

$$x = \frac{1}{a}(u-b).$$

Example 6: Compute

$$\int \frac{2x}{3x+4} \, dx.$$

The substitution

$$u = 3x + 4, \ du = 3 \, dx \implies dx = \frac{1}{3} \, du$$

can also be used to express 2x in terms of u:

$$x = \frac{1}{3}(u-4) \implies 2x = \frac{2}{3}(u-4),$$

 $\mathbf{SO}$ 

$$\int \underbrace{\frac{2}{3}(u-4)}_{u} \underbrace{\frac{1}{3}du}_{u} = \int \frac{\frac{2}{3}(u-4)}{u} \frac{1}{3}du = \frac{2}{9}\int 1 - \frac{4}{u}du$$
$$= \frac{2}{9}u - \frac{8}{9}\ln|u| + C = \frac{2}{9}(3x+4) - \frac{8}{9}\ln|3x+4| + C$$
$$= \frac{2}{3}x - \frac{8}{9}\ln|3x+4| + C$$